

A New Study of s-Process Nucleosynthesis in Massive Stars

I. The Core-Helium Burning Phase

L.-S. The,¹ M.F. El Eid,² and B.S. Meyer¹

¹Department of Physics & Astronomy, Clemson University, Clemson, SC 29634-1911

²Department of Physics, American University of Beirut (AUB), Beirut, Lebanon

ABSTRACT

We present a comprehensive study of s-process nucleosynthesis in 15, 20, 25, and 30 $M_{\odot}$ stellar models having solar-like initial composition. The stars are evolved up to ignition of central neon with a 659 species network coupled to the stellar models. In this way, the initial composition from one burning phase to another is consistently determined, especially with respect to neutron capture reactions. The aim of our calculations is to gain a full account of the s-process yield from massive stars. In the present work, we focus primarily on the s-process during central helium burning and illuminate some major uncertainties affecting the calculations. We briefly show how advanced burning can significantly affect the products of the core helium burning s-process and, in particular, can greatly deplete ^{80}Kr that was strongly overproduced in the earlier core helium burning phase; however, we leave a complete analysis of the s-process during the advanced evolutionary phases (especially in shell carbon burning) to a subsequent paper. Our results can help to constrain the yield of the s-process material from massive stars during their pre-supernova evolution.

Subject headings: nuclear reactions, nucleosynthesis, abundances –stars: evolution
–stars: interiors

1. INTRODUCTION

According to phenomenological analysis (e.g. Clayton et al. 1961, Käppeler, Beer, & Wisshack 1989), at least two distinct components are needed to fit the solar abundance distribution of the s-process nuclei which is empirically described by the well-known $\sigma N_s(A)$ curve. There is the weak s-process component which is confined to the atomic mass range $A \sim 65$ –90 (from copper to zirconium), and the main s-process component which is responsible for the production of the heavier nuclei with $A=90$ –209 (i.e. elements up to bismuth). These two components are associated with distinct stellar sites of the s-process: the weak component is mainly synthesized in massive stars ($M \geq 13 M_\odot$), while the main component is produced during recurrent helium-shell thermal pulses in AGB stars.

Our concern in this paper is a careful study of the weak s-process in massive stars. In addition to its intrinsic interest in nucleosynthesis theory, such a study is important for at least two other main reasons. First, a well determined yield from the weak component helps to constrain the contribution of the main component to the production of the nuclei in the mass range $A=65$ –90. This will help determine the degree to which s-process isotopes in mainstream SiC grains should differ from solar ratios (e.g. Gallino, Busso, & Lugaro 1997). Secondly, the weak component depends strongly on the initial metallicity (e.g Prantzos, Nomoto, & Hashimoto 1990; Baraffe, El Eid, & Prantzos 1992; Raiteri et al. 1992); thus, it may be used to study the role of massive stars in the early phase of the chemical evolution of the Galaxy.

Peters 1968 was the first to suggest that the main neutron source triggering the s-process in massive stars is the $^{22}\text{Ne}(\alpha, n)^{25}\text{Mg}$ reaction. The ^{22}Ne abundance is built up at the beginning of central helium burning by the reaction chain $^{14}\text{N}(\alpha, \gamma)^{18}\text{F}(e^+ \nu_e)^{18}\text{O}(\alpha, \gamma)^{22}\text{Ne}$, where the ^{14}N is left over from CNO cycling during the preceding hydrogen burning phase. As the temperature rises in the helium burning core, the reaction $^{22}\text{Ne}(\alpha, n)^{25}\text{Mg}$ releases neutrons which may then capture onto ^{56}Fe seed nuclei.

There are a number of key factors that influence the s-process occurring in this site:

1. The ^{25}Mg produced by $^{22}\text{Ne}(\alpha, n)^{25}\text{Mg}$ serves (along with other nuclei such as ^{12}C , ^{20}Ne , ^{16}O) as a neutron poison, thereby reducing the number of neutrons per ^{56}Fe seed.
2. The reaction $^{22}\text{Ne}(\alpha, n)^{25}\text{Mg}$ becomes effective only during the late phase of central helium burning when the core temperatures exceeds $\sim 2.5 \times 10^8$ K. Since helium ignites at temperatures less than 2×10^8 K, only a terminal fraction of the core helium burning phase actively contributes to the s-process in massive stars.
3. The production of neutrons by the ^{22}Ne source is also limited by the efficiency of the reactions competing for the alpha particles such as the $^{12}\text{C}(\alpha, \gamma)^{16}\text{O}$ reaction, which has a rate still not accurately determined to better than a factor of two, and the $^{22}\text{Ne}(\alpha, \gamma)^{26}\text{Mg}$ reaction.

4. As our computations show (see §5), s-process nucleosynthesis is rather sensitive to the treatment of mixing at the edge of the convective core during the final part of central helium burning (see discussion in sect. 5.1). The yield depends also on the mass fraction of the core that is ejected without significant additional nuclear modification. We return to this theme at the end of this paper.

Network calculations of the s-process require temperature, density, and neutron abundance histories. Due to computational limitations, past efforts to model the s-process in massive stars have relied on the technique known as “post-processing”, specifically, using temperature, density, and neutron abundance histories derived from separately calculated stellar models. This may be a reasonable treatment of the s-process during central helium burning, but for advanced burning stages (such as carbon-shell burning), rapid mixing and possible feedback effects on the nuclear energy generation rate make the post-processing approach questionable. In the calculations by Raiteri et al. 1991 and Raiteri et al. 1993, the contribution of the carbon-burning shell to the s-process in a 25 $M_{\odot}$ star was calculated on the basis of the stellar models by Nomoto & Hashimoto 1988 by assuming a convective C-shell of constant temperature and density. This is a simplified treatment in which a possible feedback effect on the stellar models and the important role of mixing are not taken into account (as we show in §6).

Given the complex behavior of the abundance pattern during the advanced evolutionary phases, our goal is to avoid using post-processing to calculate the s-process in massive stars. This is accomplished by coupling a full network of the s-process to the stellar models (see §2 and §3). In this way, we are able to follow s-process products carefully through the shell-helium burning and shell carbon burning phases in a consistent way, and thereby to gain more insight into the s-process. The results of our calculations are very rich in detail; therefore, in order to keep the present paper manageable in scope, we focus primarily on the core helium burning phase. We do, however, present some interesting results for carbon shell burning in §6.

One of our primary goals in this work is to explore the effects of key unresolved input uncertainties on the massive-star s-process yields. Differences in s-process yields between massive star models largely result from differences in the treatment of mixing within the stellar models and from the uncertainty in the $^{12}\text{C}(\alpha, \gamma)^{16}\text{O}$ rate (a factor of two) and the neutron-capture cross sections in mass region between Fe and Zr ($\sim 10\%$ uncertainty). We explore the effect of each of these uncertainties in some detail. To estimate the uncertainty and the affect of the $^{12}\text{C}(\alpha, \gamma)^{16}\text{O}$ rate on the s-process yields, we perform two calculations of a 20 $M_{\odot}$ star, one using the rate given by Caughlan & Fowler (1985) (hereafter CFHZ85) and the other one using the rate given by Caughlan et al. (1988) (hereafter CF88). We discuss the differences in the stellar models by comparing our results with those calculated by the FRANEC code (Käppeler et al. 1994; hereafter K94). Comparison of our results with those of Prantzos, Nomoto, & Hashimoto (1990; hereafter P90) allows us to consider the effect of the neutron capture cross section uncertainties.

Additional uncertainties important for the weak s-process arise from other key alpha particle

capture rates. Recent progress in eliminating these uncertainties has been made by K94 who measured reaction rates for $^{18}\text{O}(\alpha, \gamma)^{22}\text{Ne}$ and $^{22}\text{Ne}(\alpha, \gamma)^{26}\text{Mg}$. An uncertainty on the magnitude of the $^{22}\text{Ne}(\alpha, n)^{25}\text{Mg}$ reaction still remains, however. In particular, the possibility of a contribution to the rate from a low-lying resonance at 633 keV in the compound nucleus ^{26}Mg has been shown to increase the s-process production factors by an order of a magnitude above the production factors from the rate given by CF88. In order to explore the effect of this uncertainty further, we compare the s-process yields of 15, 20, 25, and 30 $M_{\odot}$ stars computed with the $^{22}\text{Ne}(\alpha, n)^{25}\text{Mg}$ rate given by CF88 and with the rate containing an additional 10% contribution from the 633 keV resonance (the WT case in K94).

Despite the uncertainties discussed above, the general notion of massive stars as the source of the weak s-process component is by now firmly in place. A few difficulties remain, however. Most significantly, a potential problem has been recognized since the work of P90. In a comprehensive study, these authors calculated s-process yields from massive stars during core helium burning as a function of stellar mass and metallicity. Feeding these as inputs into a simple chemical evolution model and comparing the integrated abundance with solar abundance, P90 concluded that massive stars during core helium burning overproduce the nucleus ^{80}Kr by a factor of $\simeq 2$, thereby presenting a serious difficulty for explaining the solar abundance of this isotope. One possible solution considered by P90 is the destruction of ^{80}Kr during the supernova explosion. This would probably only destroy abundant ^{80}Kr in the inner 2-3 solar masses. This is the matter that would for the most part remain in the stellar remnant, however, and thus the effect would be unlikely to alter the ejected ^{80}Kr abundance significantly.

Another solution of the ^{80}Kr overproduction problem has been suggested by Raiteri et al. (1991). These authors argue that the solar s-only nuclei of the weak component originate not only from massive stars but also in part from the main component (i.e. from thermal pulses in the helium shell of low-mass stars) and the p-process. In particular, Raiteri et al. 1991 find that massive stars contribute significantly to the solar system's supply of ^{80}Kr through both the s- and p-processes. Low-mass stars do not. On the other hand, they find that low-mass stars contribute abundantly to the supply of other s-only nuclei in the $A \approx 60 - 90$ region. In other words, they argue that the high production of ^{80}Kr in massive stars is needed to compensate for its relatively low production in low-mass stars. This is a plausible explanation but requires confirmation from other calculations of the s-process in both low-mass and high-mass stars.

Based on the calculations presented here and in a forthcoming paper, we find an added wrinkle to the question of large ^{80}Kr overproduction factors from massive star models. We suggest that these overproduction factors have been significantly overestimated because calculations up to the present have not yet correctly accounted for carbon-shell s-processing. Our results indicate that carbon shell burning can strongly deplete the large ^{80}Kr overabundances produced in core helium burning. Interestingly, the amount of depletion of ^{80}Kr depends strongly on the amount of ^{22}Ne left over from core helium burning as well as the extent of and the timescale for convective mixing in the carbon shell. This identifies for ^{80}Kr a potentially powerful role as a diagnostic of

advanced burning phases in massive stars. This is in addition to the important part it plays in studies of galactic abundance evolution.

The outline of the present paper is the following. In §2 we present an overview of the stellar evolution code used to obtain the stellar models. In §3, we briefly described the network we have used to calculate the s-process abundances and to determine the energy generation rates in the stellar models. In §4, §5 and §6 we summarize the results for the s-process, and in §7 present our conclusions.

2. STELLAR EVOLUTION CODE

The stellar models presented in this work were computed with a one-dimensional implicit hydrodynamical lagrangian code that solves the stellar structure equations (see Kippenhahn & Weigert 1990). The original version of this code was described in Ober, El Eid, & Fricke (1983). The present version of this code has the following general features: the discretized stellar structure equations are iterated for a given time step in terms of five variables (radius r , velocity u , temperature T , density ρ , and luminosity L). This is done for the whole stellar model from the center to the photosphere with appropriate central and surface boundary conditions. In particular, the surface boundary conditions are adjusted to a gray atmosphere in a manner similar to that described by Paczyński & Różyczka 1977. The evolutionary time step is chosen by imposing upper limits on the variations of the physical variables in the stellar model. The physical assumptions and parameterizations used to construct the stellar models are summarized as follows:

1. Rosseland mean opacities from Livermore (Iglesias & Rogers 1996) have been used for the hydrogen-rich stellar layers and for the hydrogen-free, but C/O enhanced layers. The OPAL opacities are not available for low temperatures (below about 6000 K). For such temperatures, we used the Rosseland mean opacities calculated by Alexander & Ferguson 1994 which include the effects of molecules on the opacities. Conductive opacities have also been included according to the analytical approximation by Iben 1975.
2. Partial ionizations of H, He, C, N, O, Ne, Mg and Si and their effect on the equation of state are included by solving the Saha equation taking into account all relevant excited atomic states (El Eid & Höflich 1990). In fully ionized stellar layers, the equation of state includes a full relativistic description of the degenerate electron and positron gas (Cox & Guili 1968). The effect of the Coulomb pressure on the equation of state is approximately treated according to the one-dimensional plasma calculations by Hansen 1973.
3. The extensions of the convective regions in the stellar models are determined according to the Schwarzschild criterion for convective instability, $\nabla_{\text{rad}} \geq \nabla_{\text{ad}}$. We do not include overshooting (e.g. Schaller et al. 1992; Bressan et al. 1993), or semiconvection (e.g. El Eid 1995, Langer & Maeder 1995). There is no consistent theory yet that describes these complex

effects, and their inclusion would complicate the discussion of the s-process, and prevent the comparison with the results of previous calculations. Convective energy transport is treated according to the mixing length theory (see Kippenhahn & Weigert 1990) with a mixing length parameter $l=2.0 H_P$, where H_P is the local pressure scale height.

4. Incrementing the abundances of nuclear species due to nuclear reactions is followed by integrating a detailed network of nuclear reactions as described in §3. The change of the abundances due to convective mixing is performed by solving implicitly a diffusion equation (see Langer, El Eid, & Fricke 1985; El Eid 1995) of the following form:

$$\frac{\partial X_i}{\partial t} = \frac{\partial}{\partial M_r} \left[(4\pi r^2 \rho)^2 D \frac{\partial X_i}{\partial M_r} \right], \quad (1)$$

where r is the radius, ρ is the density, M_r is the mass coordinate. The diffusion coefficient D in Eq. (1) is taken as $D = v_c l$, where v_c is the convective velocity obtained from the mixing length theory, and l is the mixing length parameter. It must be emphasized that mixing according to Eq. (1) does not always resemble instantaneous mixing. It does so only if the nuclear time scale τ_{nuc} is larger than the diffusion time scale $\tau_{\text{diff}} \sim l^2/D$. Such conditions are encountered during the early evolution phases: hydrogen, helium and carbon burning phases. Beyond the central carbon phase, however, τ_{nuc} may become comparable or smaller than τ_{diff} , especially in the case of the light particles (neutrons, protons and α -particles) which are liberated by nuclear reactions. Neutrons set free are captured before any mixing can occur.

The stellar models during these advanced phases were constructed in the following way: for a given evolution time step Δt , a stellar model is first constructed. With this time step, the network of nuclear reactions is integrated, and the resulting abundances are mixed according to Eq. (1). Subsequently, the smallest diffusion time scale τ_{diff} in the stellar model is determined. If τ_{diff} is found smaller than Δt , the network is integrated again but with a fraction of τ_{diff} (typically 1 – 10%) in order to correct the abundance profiles for all nuclear species whose lifetimes are smaller than τ_{diff} . Through a number of tests, we have convinced ourselves that this approximate method of treating time-dependent mixing leads to the correct energy generation rates in the stellar models.

5. In the present models, we have included mass loss by stellar wind according to the semi-empirical relation by De Jager et al. (1988). Mass loss for the stars in the mass range considered here has only a minor effect on the central conditions affecting the s-process nucleosynthesis during core helium burning.

Finally we mention that the neutrino energy losses due to pair, photo, plasma, bremsstrahlung and recombination processes are taken into account according to the recently up-dated analytical approximation by Itoh et al. (1996).

3. THE REACTION NETWORK

We have incorporated a large nuclear network into the stellar evolution code in order to follow the s-process nucleosynthesis occurring during advanced stages of core and shell burning. The nuclear species included in our network are those listed in table 1. Up to zinc, the network includes isotopes of each element ranging from an isotope near the proton-drip line to an isotope two mass units greater than the most massive stable isotope. This range of isotopes is needed to follow the nuclear burning through hydrostatic oxygen burning. For elements of higher charge than zinc, the isotopes included range from the lowest mass stable isotope to the isotope two mass units greater than the most massive stable isotope. This range of isotopes comfortably includes all species important for the s-process.

The required nuclear data come from a variety of sources. Strong and electromagnetic nuclear reaction rates are for the most part from CF88 and Thielemann, Arnould, & Truran 1986. The new reactions rates for $^{18}\text{O}(\alpha,\gamma)^{22}\text{Ne}$ and $^{22}\text{Ne}(\alpha,\gamma)^{26}\text{Mg}$ from K94 are applied in our calculations. The possibility that the $^{22}\text{Ne}(\alpha,n)^{25}\text{Mg}$ rate might be enhanced due to a possible resonance at 633 keV (K94) is also considered by performing test calculations including the modified rate (see §5). In the Ne-Na-Mg-Al region, we use certain updated rates from El Eid & Champagne 1995.

The rates for the neutron-capture reactions are taken from Beer, Voss, & Winters (1992). These authors have considered the dependence of these cross sections on temperature. If the s-process occurs at high temperature, for example near $kT=90$ keV as in carbon-shell burning in massive stars, then the neutron-capture cross sections no longer follow the simple $1/v$ -law and the $kT=30$ keV thermally-averaged cross sections (Bao & Käppeler 1987) are no longer strictly valid. Therefore, we have used the Beer, Voss, & Winters (1992) cross sections to permit study of the s-process under relatively extreme conditions.

The nuclear masses and most weak interaction (electron-capture and β -decay) rates are taken from Tuli 1995. Certain weak rates are temperature and density dependent. For these rates we use the calculations of Takahashi & Yokoi 1987. Extrapolations of some weak interaction rates (such as the β -decay rate of ^{79}Se) to higher temperatures were necessary and were done based on experimental results (Klay & Käppeler (1988) and also the data given in Table 2 of Raiteri et al. 1993.

4. CHARACTERISTICS OF THE EVOLUTIONARY MODELS

We have evolved models of stars of masses 15, 20, 25, and $30 M_{\odot}$ with initial solar-like composition (see Anders & Grevesse (1989)) from the main sequence up to the ignition of central neon. The network described in §3 has been coupled (between the time steps) to the stellar models during all these evolutionary phases. In this way, we are able to determine correctly the neutron density during all these burning phases. To our knowledge, no such extended calculations have

been done before for the massive stars considered here. For example, in the recent work by Chieffi et al. 1998 the effect of the neutron capture reactions has been ignored during central helium burning, with the result that many isotopic abundances which are affected by neutron capture reactions may not be well determined.

In total, we ran ten stellar models. These are summarized in Table 2. Our reference calculations are case A, for which we used a reaction rate expression for $^{12}\text{C}(\alpha, \gamma)^{16}\text{O}$ 1.7 times larger than the CF88 value (as suggested by the work of Weaver & Woosley 1993) and the CF88 rate expression for $^{22}\text{Ne}(\alpha, n)^{25}\text{Mg}$. The case B calculations were identical to those in case A except we included a low-energy resonance in the $^{22}\text{Ne}(\alpha, n)^{25}\text{Mg}$ reaction. In particular, we used the rate expression from K94 which included a 10% contribution from this resonance. For both case A and B we computed models for 15, 20, 25, and 30 $M_{\odot}$ stars. To explore the sensitivity of our results to the $^{12}\text{C}(\alpha, \gamma)^{16}\text{O}$ reaction, we also ran 20 $M_{\odot}$ models for the CF88 (case C) and CFHZ85 values for the reaction rate (case D). We find interesting consequences for both of these rates.

In the following, we describe some general properties of our stellar models. We first present in Fig. 1 the evolution of the central temperature T_c versus the central density during the various evolutionary phases indicated there for the four case A stars. Calculating the stars through these phases means that we can also follow the s-process in regions where the different shell sources are active, especially the contribution of the carbon-burning shell to the s-process. The dependence of the central temperature T_c on the central density ρ_c (in the phases before central carbon ignition) is reasonably well approximated as $T_c \approx 2 \times 10^7 (\rho_c [\text{g cm}^{-3}])^{1/3}$ K, although in general the more massive stars reach given central temperatures at lower densities or have higher central temperatures for given densities. The $T_c \propto \rho_c^{1/3}$ indicates cores of our massive star models behave approximately as polytropes of index three, showing that the radiation pressure in the stars is not negligible (e.g. Clayton 1983). After the central carbon ignition, the $T_c \propto \rho_c^{1/3}$ relation is modified as neutrino energy loss radiates away entropy from the core (e.g. Arnett 1996).

Table 3 we summarize some of the details of our four case A stellar models at key stages in their evolution. Our stellar models end their central helium burning as red supergiants. Also interesting is the increase of the helium core mass M_{α} with time, which indicates the efficiency of accretion from the hydrogen-shell burning. After the end of core carbon burning or even somewhat earlier, however, M_{α} is essentially fixed because of the short duration of the subsequent burning phases. An MPEG movie showing the evolution of temperatures, densities, luminosities, and mass fractions of ^1H , ^4He , ^{12}C , ^{16}O , and ^{20}Ne as a function of internal mass radius in a 25 $M_{\odot}$ star can be viewed at the Clemson nuclear astrophysics web site, <http://photon.phys.clemson.edu/movies.html>.

Table 4 shows the overabundances inside the hydrogen exhausted core at the end of core hydrogen burning for the four case A models (stage 2 in table 3). Only those stable or long-lived isotopes with overabundances noticeably different from unity are shown. These are the isotopes that changed during hydrogen burning. Because the stellar models all began with

solar composition, overabundances of unity equate to solar abundances. Some interesting changes have occurred during the core hydrogen burning. ^1H , ^3He , and ^7Li are all strongly depleted, as expected. ^4He is enhanced by production from hydrogen. The abundances of the CNO isotopes have shifted around giving enhanced ^{13}C and ^{14}N , as expected. Ne-Na-Mg cycle burning alters the abundances of isotopes of fluorine to magnesium. The neutron source ^{22}Ne is depleted at this stage. It will be created during subsequent He burning. Finally, the abundances of ^{138}La , ^{163}Dy , and ^{187}Re all drop due to temperature enhanced weak-decay rates. The abundance of ^{187}Os rises because it is the daughter of ^{187}Re decay. The abundances of all other isotopes, however, are unaffected by the core hydrogen burning.

The next stage of the star’s evolution, and the one of primary interest for the present paper, is core helium burning. The evolution of each convective core mass M_{cc} for the stars under study during central helium burning is shown versus time in Fig. 2 (upper panel), and versus central helium mass fraction, $X(^4\text{He})_c$ (referred to as Y_c in the following) in Fig. 2 (lower panel). The step-like structures visible in these figures can be easily removed by finely resolving the edge of the convective core, that is, by adding more meshpoints there. Our experience shows that this kind of fine resolving is not required as long as Y_c is larger than 0.10, since the convective core grows sufficiently slowly that a sudden uncontrolled injection of helium into the core does not occur. When Y_c decreases below 0.10, however, the above stable situation changes drastically. As the helium supply dwindles, the core contracts gravitationally. The convective core grows in response to the release of the gravitational binding energy. This allows the helium-depleted core to ingest suddenly a large amount of the overlying ^4He which in turn dramatically increases the central energy generation rate. This halts the central contraction and allows the convective core to shrink again. This behavior is called “breathing pulses” in the literature (e.g. Castellani et al. 1985). We think that the breathing pulses are not physical. They are rather a numerical artifact of the necessity of discretizing the stellar model. They can be avoided by careful rezoning at the edge of the convective core, or perhaps also by coupling mixing and nucleosynthesis during the iteration cycle, which is of course extremely time consuming. We choose the former strategy in the present calculations with the result that the convective core grows smoothly as seen in Fig. 2 below $X(^4\text{He})=0.10$ and thus that the helium mixed into the convective core during each time step behaves in a regular way.

To make the nuclear consequences of this point more clear, we note that without the fine resolution of the convective core edge but also without any limitation on the breathing pulses we would not find the relatively high mass fractions of ^{12}C at the end of central helium burning given in Table 5. For example, had we not finely resolved the edge of the convective core in our case A $25\text{ M}_\odot$ stellar model, the resulting central ^{12}C and ^{16}O mass fractions would have been 0.17 and 0.79, respectively, in place of the corresponding values 0.22 and 0.75 in case A. The breathing pulses allow sudden growth of the convective core and ingestion of additional ^4He and, consequently, more conversion of ^{12}C into ^{16}O via the $^{12}\text{C}(\alpha, \gamma)^{16}\text{O}$ reaction. Importantly for the s-process, the breathing pulses also burn more ^{22}Ne and yield a more robust s-process (e.g. K94).

Also consequential is the fact that the greater the destruction of ^{22}Ne in core helium burning, the less efficient will be the s-process in later burning stages, particularly shell helium and carbon burning.

Stellar modelers typically avoid the problem of breathing pulses by “limiting” the growth of the convective core at low core helium mass fractions, that is, by using computer code commands to prevent the core from growing in size once the central helium mass fraction falls below some value. We ran another $25\text{ M}_{\odot}$ calculation identical to our case A model but without fine mass resolution of the convective core edge and including a limitation on the growth of the core for $Y_c < 0.10$. The resulting central ^{12}C and ^{16}O mass fractions were 0.25 and 0.73, respectively. The limiting prevented any core growth at low central helium and thus allowed less ^{12}C to burn to ^{16}O . For the same reason, less ^{22}Ne burned, thereby giving a weaker s-process.

The model with suppression of the breathing pulses but without fine mass resolution gives results reasonably close to our case A model. Nevertheless, the case A model is clearly the more satisfactory of the two because it does not have an artificial constraint on it. We have therefore chosen in our models to resolve the edge of the convective core finely since it is most realistic scenario for our chosen convection criterion and gives the most believable results for the s-process.

In Table 6 we compare the details of core helium burning in our four case A stars with the results of other authors, where available. In general, our results agree well with those of P90, who used the stellar models of Nomoto & Hashimoto 1988, both in terms of the maximum convective core mass and the helium burning lifetime. This makes sense because both our models and those of Nomoto & Hashimoto 1988 used the Schwarzschild criterion for convection and employed no overshooting. On the other hand, both sets of models differ from those computed with the FRANEC code (K94). In particular, our maximum convective core masses are at least $\sim 10\%$ smaller than those produced with FRANEC. At the same time, the helium burning lifetimes in the FRANEC models are considerably shorter than in our models and those of P90. The larger cores maintain a higher supply of ^4He and a faster ^4He consumption rate. It is interesting that the FRANEC calculations presented in K94 are rather similar to those in Schaller et al. 1992, which included moderate overshooting (by 20% of the pressure scale height). It is eventually the treatment of mixing at the edge of the convective core that simulates the effect of overshooting, even though it is probably not explicitly included in the FRANEC calculations. These stellar model differences have implications for s-process nucleosynthesis, as discussed §5.

5. s-PROCESS DURING CENTRAL HELIUM BURNING

In this section we present and analyze our results for the s-process. We also compare our findings with those of other authors.

5.1. s-Process Diagnostics

Two quantities usefully characterize the efficiency of s-process nucleosynthesis. One is the number of neutrons captured per iron seed:

$$n_c \equiv \sum_{A=56}^{209} (Y_A - Y_A^0)(A - 56)/Y_A^0, \quad (2)$$

where Y_A is the final abundance of a nuclear species of mass number A , and Y_A^0 is its initial value. The other quantity is the neutron exposure defined as:

$$\tau_n(m_r) \equiv \int_0^t n_n(m_r, t') v_{th} dt', \quad (3)$$

where n_n is the neutron density at a relative mass m_r , and $v_{th} = (2kT/m_n)^{1/2}$ is the thermal velocity of the neutrons. In the present calculations, we take v_{th} evaluated at $kT = 30$ keV.

It is possible to define an average neutron exposure according to the above definition if the s-process occurs in a convective core as during central helium burning in massive stars, that is

$$\langle \tau \rangle = \int_0^t \langle n_n \rangle v_{th} dt', \quad (4)$$

where $\langle n_n \rangle$ is the neutron density averaged over the convective core and v_{th} is the neutron thermal velocity at 30 keV. However, such a quantity represents only a rough indicator of the efficiency of the s-process, since the neutrons are actually locally absorbed, mostly at the central region of the star. A more reliable quantity is an exposure defined in terms of a nucleus like ^{54}Fe which is only destroyed during the s-process:

$$\tau_{54} = -\frac{1}{\sigma_T} \ln(X_{54}/X_{54}^0), \quad (5)$$

where σ_T is the neutron-capture cross section at some temperature T (usually taken to be 30 keV), and X_{54} , X_{54}^0 are the final and initial mass fractions of that nucleus respectively.

5.2. Our Results

Figs. 3a-d show the central τ_n , τ_{54} , n_c , n_n and $X_i/X_{i\odot}$ versus central $X(^4\text{He})$ for a $25 M_\odot$ stellar model in cases A and B. Again, case A is the reference calculation which uses the CF88 rate for the $^{22}\text{Ne}(\alpha, n)^{25}\text{Mg}$ reaction, while case B includes a 10% contribution from a putative resonance. According to the discussion by K94, the contribution of such a resonance to this reaction rate cannot yet be excluded on experimental grounds.

From Figs. 3a-d it is clear that the effect of the resonance would be to augment the s-process. For example, Fig. 3b shows that the central ^{22}Ne abundance decreases much more dramatically in

case B than case A. The result is a larger release of neutrons and, consequently, a higher n_c (Fig. 3c), considerably more ^{80}Kr and ^{25}Mg (Fig. 3b), and a larger central τ_n (Fig. 3a). Interestingly, the sharp increase of the ^{80}Kr mass fraction occurs earlier in time (or at higher Y_c) in case B than in case A due to the more efficient ^{22}Ne destruction. The 15, 20, and 30 $M_\odot$ models show qualitatively similar results; therefore, on the basis of these results, we can conclude that a tight constraint on the s-process in massive stars is not possible as long as the rate of $^{22}\text{Ne}(\alpha, n)$ reaction is experimentally unsettled.

Fig. 3a shows the two neutron exposures τ_n and τ_{54} . It is important to distinguish them. The τ_n shown in Fig. 3a is the neutron exposure at the center of the star, that is, the τ_n computed from eq. (3) at $m_r = 0$. It is thus the neutron exposure that would have been seen by nuclei that always remained in the center of the star. The value of this quantity at the end of core helium burning is greater than about 4 mb^{-1} , which is an extraordinarily large exposure. It results from the high neutron density in the core of the star, as shown in Fig. 4, where the temperature is highest and the ^{22}Ne burns most effectively. Outside the very core of the star, the temperature is lower and the neutron density is considerably less.

A neutron exposure of 4 mb^{-1} or greater is large enough to drive all initial iron seed nuclei up to bismuth. This, of course, is not what happens. Because of convective mixing, a given nucleus spends only a small fraction of the helium-burning time in the star’s center. As mentioned above, the neutron density outside the very central region of the star is much less, so little s-processing is occurring there. This is illustrated in Fig. 5, which shows τ_n as a function of interior mass for our eight models. Only the inner few tenths of a solar mass have a significantly large τ_n , and it is only here that neutron captures are occurring. Nuclei are convected down into these inner regions, capture neutrons, and then mix back out into the core. The result is that nuclei, on average, see an exposure that is much less than the τ_n at the center.

A better measure of the average neutron exposure seen by nuclei during core helium burning is τ_{54} at the center of the star. It is shown in Fig. 3a and is instantaneously computed from eq. (5) using the current value of the ^{54}Fe mass fraction. The definition of the mass fraction of ^{54}Fe , $X(^{54}\text{Fe})$, is unambiguous because $X(^{54}\text{Fe})$ is uniform throughout the convective zone. ^{54}Fe nuclei, which are only destroyed during the s-process, are constantly capturing neutrons in the center of the star and mixing throughout the core; thus, τ_{54} is an appropriate measure of the average neutron exposure. The finding is that this average exposure is considerably less than τ_n at the center of the star, as expected, and better characterizes the resulting s-process abundances. Nevertheless, even τ_{54} is not the complete picture. A single s-process abundance pattern characterized by the final neutron exposure τ_{54} is not an adequate description of the true abundance pattern at the end of core helium burning. In fact, a distribution of neutron exposures is needed to characterize the final abundances. The reasons for this are complex, and we address them in a forthcoming paper.

To illustrate the effect of the $^{12}\text{C}(\alpha, \gamma)^{16}\text{O}$ reaction rate on the s-process, we present Fig. 6.

This figure shows the results for the s-process for the 20 $M_{\odot}$ star in cases A, C, and D. For the lowest value for the reaction rate (case C: CF88), the s-process is considerably more robust than for higher values of the reaction rate. The reason, as pointed out by Raiteri et al. 1991, is that alpha particle capture on ^{12}C competes with that on ^{22}Ne for consumption of ^4He ; thus, a lower rate of $^{12}\text{C}(\alpha, \gamma)^{16}\text{O}$ leaves more alpha particles available for the $^{22}\text{Ne}(\alpha, n)^{25}\text{Mg}$ reaction and consequently leads to more neutron production. It aids understanding to recall that most of the ^{22}Ne consumption occurs when less than 10% of the helium remains (Fig. 3b and 6b).

The faster $^{12}\text{C}(\alpha, \gamma)^{16}\text{O}$ reaction rate explains the less efficient s-process in case A than in cases C and D. Counter to expectations, however, case D shows a more efficient s-process than case A, even though the $^{12}\text{C}(\alpha, \gamma)^{16}\text{O}$ reaction rate is always greater in case D over the temperature range of interest for the core helium. The resolution of this puzzle lies in the fact that the faster $^{12}\text{C}(\alpha, \gamma)^{16}\text{O}$ reaction rate (Case D rate is $\simeq 2.4 \times$ rate of Case C at temperature $\sim 2.5 \times 10^8$ K) also leaves fewer ^{12}C nuclei around (^{12}C mass fraction at the end of core He burning is 0.12 in Case D and 0.23 in Case C). The smaller abundance of ^{12}C can compensate the faster rate leading to a larger abundance of alpha particles and greater consumption of ^{22}Ne . When the triple-alpha reaction is no longer the major consumer of alpha particles (owing to the small value of Y_c), the ^{12}C and ^{22}Ne compete in proportion to their abundances for the final helium. In fact, careful examination of Fig. 6 shows that the various curves behave in strict accordance with the rule that a larger $^{12}\text{C}(\alpha, \gamma)^{16}\text{O}$ reaction cross section gives less s-processing until the ^4He mass fraction drops to about 0.01. Then cases A and D cross over. It is here that the lower abundance of ^{12}C in case D overcomes the larger reaction cross section.

The same interplay between ^{12}C and ^{22}Ne explains the difference between the case A and B central ^{12}C mass fractions in table 5. One would expect a faster $^{22}\text{Ne}(\alpha, n)^{25}\text{Mg}$ rate (case B) to allow ^{22}Ne to compete more effectively for alpha particles, thereby resulting in a higher final ^{12}C mass fraction. In fact, the opposite is true. The case B ^{12}C mass fractions are consistently slightly *lower* in table 5 than the corresponding case A values. The reason is that so much ^{22}Ne burns in the case B models that its low abundance overcomes the faster rate and makes it compete for alpha particles slightly less effectively than in case A.

5.3. Comparison of s-process Studies

In Table 7 we present a comparison of our results for the s-process at helium exhaustion and those obtained by P90 and K94 when all three sets of calculations used the CF88 expression for the $^{22}\text{Ne}(\alpha, n)^{25}\text{Mg}$ reaction rate. Before discussing the differences, it is useful to point out two consistent trends within the results of each set of authors. First, the more massive the star, the more ^{22}Ne is burned. This reflects the higher core temperatures in more massive stars and, consequently, the higher rate of alpha particle capture on ^{22}Ne . Second, it is consistently the case that the lower the ^{22}Ne abundance at core helium exhaustion, the greater the number of neutrons captured per iron seed nucleus. This is simply due to the fact that the neutrons driving

the s-process are liberated from ^{22}Ne .

Now we point out the differences between our results and those of the other authors. First, (using our reference calculation case A), our models typically burn less ^{22}Ne than P90 but more than K94. Second, we consistently get more s-processing than either P90 or K94.

Our models typically burn less ^{22}Ne than P90 because these latter authors used the CF88 rate value for the $^{12}\text{C}(\alpha, \gamma)^{16}\text{O}$ reaction. As with our case C model discussed in the previous subsection, a low value for this rate means more alpha particles are available for capture on ^{22}Ne during the s-processing phase.

Despite the fact that our models typically burn less ^{22}Ne than those of P90, our models in fact show a larger number of neutrons captured per iron seed nucleus. We believe the reason for this is that our calculations used temperature-dependent neutron capture cross sections (Beer, Voss, & Winters (1992)) while P90 used the 30 keV cross sections of Bao & Käppeler 1987. Much of the s-processing in our models happens at temperatures in the range of about 23 to 26 keV, and the cross sections for typical s-process nuclei in this temperature range can be up to factors of two larger than at 30 keV. On the other hand, the cross sections for major s-process poisons at 23 to 26 keV can be slightly lower than at 30 keV. Our models therefore also show greater s-processing because of more efficient neutron capture by heavy nuclei. It is of interest to note that the rather large differences in the capture cross sections as a function of temperature probably reflect the role of resonances in the compound nuclei.

Finally, it is worth pointing out that our case C $20\text{ M}_{\odot}$ stellar model leaves less ^{22}Ne than the corresponding P90 model even though they both used the same $^{12}\text{C}(\alpha, \gamma)^{16}\text{O}$ reaction rate value. Our best explanation for this is some difference in the stellar models, perhaps in differing convective histories. This is plausible since there are rather strong differences in $\langle \tau \rangle$ between our models and those of P90. Interestingly, our case D $20\text{ M}_{\odot}$ model burns about the same amount of ^{22}Ne as does the corresponding P90 $20\text{ M}_{\odot}$ model. Our model shows more s-processing, however, which we again attribute to somewhat higher neutron capture cross sections in our models during the s-process phase.

The models of K94 show consistently less s-processing than either our models or those of P90. This is true despite the fact that our models compare rather favorably to those of K94 in terms of $\langle \tau \rangle$ and maximum neutron density averaged over the convective core. These differences are also present despite the fact that the neutron capture cross sections used in the two sets of calculations are the same and the $^{12}\text{C}(\alpha, \gamma)^{16}\text{O}$ reaction rates do not differ greatly. To explain the differences, therefore, we again appeal to differences in stellar models. Table 6 compares the maximum convective core masses and helium burning durations of the various models. Our models compare quite well with those of P90. On the other hand, the models of the FRANEC code used in K94 have consistently larger maximum convective cores and shorter helium burning lifetimes. Interestingly, the convective cores and helium burning durations of the K94 models are closer to those of Schaller et al. 1992 which included convective overshooting during core helium burning.

The treatment of mixing in the FRANEC models presented by K94 leads to enlarged convective core masses similar to the overshooting case. Whatever the reason for the differences in the stellar models, the shorter helium burning lifetimes in the K94 allowed for less s-processing compared to our models and those of P90.

In Table 8 we present a comparison of s-process results when the tentative $^{22}\text{Ne}(\alpha, n)^{25}\text{Mg}$ rate of K94 is adopted. The first obvious conclusion is that the s-processing is considerably stronger both in our models and those of K94 because of the larger rate for alpha capture on ^{22}Ne . We consistently find a larger number of neutron captures per iron seed nucleus, however, again most likely due to differences in stellar models, as discussed above. It is interesting that we find smaller overproductions of ^{80}Kr in the 25 and 30 $M_{\odot}$ stellar models than do K94. This is due to the fact that the s-process in our models is so strong that enough neutron capture has occurred to push the peak overproductions past Kr and into the Sr region (see the next subsection). For the same stellar masses, the models of K94 have overproductions still peaking at ^{80}Kr .

5.4. s-Process Production Factors

The production factors at the end of central helium burning are displayed in Fig. 7 and 8 and the numerical values are presented in Table 9. The results show the well known feature of the weak s-process component that is the synthesis of the heavy elements in the mass range $A=60-90$, with the largest production factor for ^{80}Kr .

An interesting result is that in our models with the $^{22}\text{Ne}(\alpha, n)^{25}\text{Mg}$ rate of K94 (case B), ^{80}Kr is no longer the most overproduced isotope in core helium burning in the 20, 25, and 30 $M_{\odot}$ stellar models. ^{86}Sr or ^{87}Sr show larger overproductions. These isotopes are factors of two or three less overproduced when the $^{22}\text{Ne}(\alpha, n)^{25}\text{Mg}$ rate of CF88 is used.

Apart from the large overabundances of the weak s-process nuclei, table 9 shows significant production of several interesting isotopes. ^{12}C and ^{16}O are produced as the main products of helium burning. Considerable ^{22}Ne is present because it was left unburnt. $^{25,26}\text{Mg}$ are produced via alpha particle capture on ^{22}Ne . A number of isotopes in the mass range 30-60 are made by neutron capture. Chief among these is ^{40}K . Above mass number 104, the only significant production is of ^{152}Gd , produced by branching at ^{151}Sm and ^{152}Eu , and of ^{180}Ta . This latter isotope is made in the mechanism first described by Yokoi & Takahashi 1983. The idea is that ^{179}Ta is unstable to electron capture in the star, but temperature-enhanced β^- decay from ^{179}Hf in the helium-burning core allows a steady-state abundance to build up and neutron capture to ^{180}Ta . We mention that in our current models we allow no equilibration between the ^{180}Ta ground state (which decays to ^{180}Hf in 8.15 hours) and the metastable state at 75 keV (which is essentially stable). The large spin difference between the ground and metastable states prevents direct transitions between them; thus, equilibration would have to occur through higher-lying nuclear levels. This could occur at the temperatures present in core helium burning through

intermediate states at energies in the range of 500 - 1000 keV. Were such states indeed present, the metastable state could depopulate, thereby resulting in less ^{180}Ta production. The experimental situation concerning the existence of such states is not yet clear. One group finds evidence for the existence of at least one such state (Schegel et al. 1994; von Neumann-Cosel et al. 1997) while another does not (Schumann et al. 1997). A firm prediction of ^{180}Ta production in any s-process environment will require clarification of the nuclear physics in this interesting region of the chart of the nuclides.

A final comment concerns the mainstream presolar SiC grains, which, having condensed in the outflow from AGB stars, the site of the main component, should contain differences from solar abundances that compensate for the portions produced in massive stars. An interesting puzzle is the fact that the Sr isotopes are almost normal in SiC grains (Nicolussi et al. 1998; Podosek 1998), whereas our calculations of the massive-star component show large ratios for $^{86}\text{Sr}/^{88}\text{Sr}$ and $^{87}\text{Sr}/^{88}\text{Sr}$. This imbalance ultimately requires explanation either through Galactic abundance evolution of the Sr isotopes or perhaps their modification in more advanced stellar burning phases.

6. The s-Process in Advanced Burning Phases: A Preview

Though our primary purpose in this paper has been to study the s-process during core helium burning, we briefly present in this section some results of the s-process in advanced burning phases as a preview to a more extended study to appear in a forthcoming paper. We summarize some of the most interesting results for the 25 $M_{\odot}$ model in Fig. 9. The upper panel of this figure shows τ_n and τ_{54} as functions of interior mass at the end of core helium burning (dashed lines) and near the end of core neon burning (solid lines). The peaks in τ_n at $m_r \approx 1.6 M_{\odot}$ and $m_r \approx 5.8 M_{\odot}$ indicate regions of strong neutron capture in the carbon burning and helium burning shells, respectively. The s-processing during these phases, resulting largely from burning of residual ^{22}Ne , increases τ_{54} throughout the star over its value at the end of core helium burning.

The s-processing occurring during these shell burning phases has a dramatic effect on the overabundance of ^{80}Kr . As seen in the middle panel of the figure, core and shell carbon burning deplete by factors of ten or more the huge ^{80}Kr overabundance left from core helium burning. The destruction occurs because of the high neutron densities (up to several times 10^{10} cm^{-3}) present in the innermost part of the carbon-burning shell. Because of the high neutron density in this zone, ^{79}Se captures a neutron to ^{80}Se faster than it beta decays. This in turn allows the s-process flow for the most part to bypass ^{80}Kr , thereby depleting its previously large abundance. The intense energy release due to the carbon burning in this zone drives convection out to 5 $M_{\odot}$; therefore, nuclei mix down into the high-neutron-density zone, capture neutrons, and mix back out into the outer shell. This mixing and burning depletes the ^{80}Kr throughout the convective shell. Significantly, ^{22}Ne is also continually mixing down into the burning zone. This replenishes the neutron source and keeps the neutron density high for a much longer time than would be the case without mixing. An MPEG movie illustrating the evolution of the overabundances of ^{80}Kr ,

^{22}Ne , ^{13}C , ^4He and neutron density in the $25\text{ M}_\odot$ stellar model is available for viewing on the world-wide web at <http://photon.phys.clemson.edu/movies.html>.

It is interesting that destruction of ^{80}Kr constrains the mixing in the carbon shell to be very rapid. The lifetime of a ^{79}Se nucleus against beta decay is only about 0.4 years in the innermost part of the shell at $1.6\text{ M}_\odot$ (where the temperature is about 10^9K) and about 4 years in the outermost part of the shell at $5\text{ M}_\odot$ (where the temperature is about $3 \times 10^8\text{K}$). In order to ensure that ^{79}Se nuclei throughout the shell can be re-exposed to neutrons before they beta decay and that ^{80}Kr in fact gets bypassed, the timescale for overturn of the carbon shell must be less than several tenths of a year. Our stellar models give such rapid mixing. The convective velocities in the carbon-burning shell are of the order of 1 km/s while the physical size of the shell is $\sim 40,000\text{ km}$ thick. The shell overturns once every $\sim 10^5\text{ s}$ or 0.03 yr , sufficiently fast to re-expose the ^{79}Se nuclei to the intense flux.

Interestingly, ^{80}Kr is also destroyed outside about $5.2\text{ M}_\odot$ due to shell helium burning, which yields neutron densities as high as a few times 10^{10} cm^{-3} at times earlier than that shown here. The neutron density is comparable to that in the carbon burning shell. This is somewhat surprising because the temperature and thus the the $^{22}\text{Ne}(\alpha, n)^{25}\text{Mg}$ cross section are both lower in the helium burning shell. On the other hand, the alpha particle abundance is considerably higher, allowing for efficient production of neutrons. Because of the efficient burning in the He and C shells, large ^{80}Kr overproductions from core helium burning only persist in a thin region between the outer edge of the convective carbon-burning shell and the inner edge of the helium-burning shell in our models. The lower panel of the figure shows the abundances of the major species at the given moment in the star's life.

In summary, our results indicate that ^{80}Kr , vastly overproduced in core helium burning, can be subsequently largely destroyed in advanced burning phases. The reduced yield of ^{80}Kr due to destruction during advanced burning phases has strong implications for the chemical evolution of that isotope. We intend to analyze this question in much more detail in a forthcoming paper.

7. CONCLUSIONS

We have performed detailed calculations of the s-process in massive stars with up-to-date neutron capture cross sections (Beer, Voss, & Winters (1992)), temperature and density-dependent weak interaction rates (Takahashi & Yokoi 1987; Klay & Käppeler (1988)), and a full s-process reaction network coupled to the stellar model. Consequently, we can determine the composition after core helium burning phase more accurately than calculations using post-processing. In particular, we are able to determine the dependence of the neutron exposure (τ) and neutron capture per iron seed nucleus (n_C) on mass radius at each time step, especially for the advanced evolution phases where average values are no longer possible or relevant.

From the results presented in this paper, we conclude

1. Fine rezoning of the the mesh zones near the edge of convective core can avoid the so-called core breathing problem during helium burning. The convective cores in our models grow and shrink smoothly; therefore, our treatment of the convective mixing is accurate within the framework of the Schwarzschild criterion.
2. A possible low-energy 633 keV resonance in the $^{22}\text{Ne}(\alpha, n)^{25}\text{Mg}$ reaction, which would dramatically increase the rate over the CF88 value, will significantly enhance the robustness of the s-process in massive stars. This confirms the results of K94. Accurate constraints from massive star s-processing must therefore await experimental resolution of the question of the existence of this resonance.
3. The $^{12}\text{C}(\alpha, \gamma)^{16}\text{O}$ reaction has an important effect on the efficiency of the s-process in massive stars. In particular, a low value for the rate of this reaction (such as that given in the compilation of CF88) leads to a more robust s-process than a high value (such as that of CFHZ85). The reason is that the slower this rate is, the more alpha particles are available for capture on ^{22}Ne . Nevertheless, two popular parameterizations for this rate (the CFHZ85 rate and 1.7 times the CF88 rate) give similar s-process results to within $\sim 10\text{-}20\%$.
4. Neutron captures in the core-helium-burning s-process occur only in the most central regions of the massive star because this is where ^{22}Ne is burning and releasing neutrons. For this reason, use of $\langle \tau \rangle$, the average exposure over the convective core is not a particularly useful characterization of the efficiency of the s-process. A much more useful quantity is τ_{54} which accounts for highly-concentrated burning in the center of the star and for convective mixing. Nevertheless, even this quantity is limited in its usefulness because there is in fact a distribution of neutron exposures. We will address this issue in a forthcoming paper.
5. The massive star s-process is highly sensitive to the size of the convective core and the treatment of mixing at its edge. For example, extra mixing (such as overshooting) can dramatically change the duration of the helium burning phase and, consequently, the efficiency of the s-process (see §5). Once nuclear physics issues such as the correct value of

the $^{22}\text{Ne}(\alpha, n)^{25}\text{Mg}$ reaction rate are settled, the s-process will be a powerful diagnostic of convection in massive stars.

6. The high ^{80}Kr overproduction built up during core helium burning may subsequently be strongly destroyed during carbon shell burning (§6). We will explore this possibility in a forthcoming paper.

We are grateful to Donald Clayton for helpful comments and suggestions. This work has been supported by NASA grants NAG5-4329, NAGW-3277, and NAGW-3480 and by a UR grant (5-30-1911-42-0029) at Clemson University.

REFERENCES

- Alexander, D.R., & Ferguson, J.W., 1994, *ApJ*, 437, 879
- Anders, E., & Grevesse, N. 1989, *Geochim. Cosmochim. Acta* 53, 197
- Arnett, W. D. 1996, *Supernovae and Nucleosynthesis*, (Princeton: Princeton University Press)
- Arnett, W.D., & Thielemann, F.-K. 1985, *ApJ*, 295, 589
- Baraffe, I., & El Eid, M.F. 1991, *A&A*, 245, 548
- Baraffe, I., El Eid, M.F., & Prantzos, N. 1992, *A&A*, 258, 357
- Bao, Z. Y., & Käppeler, F. 1987, *Atomic Data Nucl. Data Tables* 36, 411
- Beer, H., Voss, F., & Winters, R. R. 1992, *ApJS*, 80, 403
- Bressan, F. Fagotto, G., Bertelli, G., & Chiosi, C. 1993, *A&AS* 100, 674
- Castellani, V., Chieffi, A., Pulone, L., & Tornambè, A. 1985, *ApJ*, 296, 204
- Caughlan, G. R., & Fowler, W. A. 1988, *Atomic Data Nucl. Data Tables* 40, 283 (CF88)
- Caughlan, G. R., Fowler, W. A., Harris, M., & Zimmermann, B. 1985, *Atomic Data Nucl. Data Tables* 40, 283 (CFHZ85)
- Chieffi, A., Limongi, M., & Straniero O. 1998, *ApJ*, 502, 737
- Clayton, D. D., 1983, *Principles of Stellar Evolution and Nucleosynthesis*, (Chicago: The University of Chicago Press)
- Clayton, D. D., Fowler, W. A., Hull, T. E., & Zimmerman, B. A. 1961, *Ann. Phys.*, 12, 331
- Cox, J.P., & Guili, R.T. 1968, *Principle of Stellar Evolution*, Vol. II, (New York: Gordon and Breach)
- de Jager, C., Nieuwenhuijzen H., Van der Hucht, K. 1988, *A&AS*, 72, 259
- El Eid, M.F. 1995, *MNRAS*, 275, 983
- El Eid, M.F., & Champagne, A. 1995, *ApJ*, 451, 298
- El Eid, M.F., & Höflich, P., 1990, in *Proc. 6th Workshop on Nuclear Astrophys.*, ed. W. Hillebrandt, E. Müller, MPA, Garching, 96
- Gallino, R., Busso, M., & Lugaro, M. 1997, in *Astrophysical Implications of the Laboratory Study of Presolar Materials*, ed. T. J. Bernatowicz, E. Zinner, (Woodbury, NY: AIP), 115
- Hansen, J.P. 1973, *Phys. Rev.*, A8, 3096

- Iben, I. Jr. 1975, *ApJ*, 196, 525
- Iglesias, C.A., & Rogers, F.J. 1996, *ApJ*, 464, 943
- Itoh, N., Hayashi, H., & Nishikawa, A. 1996, *ApJS*, 102, 411
- Käppeler, F., et al. 1994, *ApJ*, 437, 396 (K94)
- Käppeler, F., Beer, H., & Wisshack, K. 1989, *Rep. Prog. Phys.*, 52, 945
- Klay, N., & Käppeler, F. 1988, *Phys. Rev. C*, 38, 295
- Kippenhahn, R., & Weigert, A. 1990, *Stellar Structure and Evolution*, (Berlin:Springer Verlag)
- Lamb, S. A., Howard, W. M., Truran, J. W., & Iben, I., Jr. 1977, *ApJ*, 217, 213
- Langer, N., & Maeder, A., 1995, *A&A*, 295, 685
- Langer, N., El Eid, M.F., & Fricke, F. 1985, *A&A*, 145, 179
- Nicolussi G. K., Pellin M. J., Lewis R. S., Davis A. M., Clayton R. N., & Amari S. 1998, *PRL*, 81, 3583
- Nomoto, K., & Hashimoto, M. 1988, *Phys. Rep.* 163, 13
- Ober, W., El Eid, M.F., & Fricke, K.J. 1983, *A&A*, 119, 61
- Paczyński, B., & Różyczka, M. 1977, *Acta Astron.*, 27, 214
- Peters, J.G. 1968, *ApJ*, 154, 224
- Podosek, F., Prombo, C.A., Amari, S., & Lewis, R.S. 1998, *ApJ*, in press
- Prantzos, N., Arnould, M., & Arcoragi, J.-P. 1987, *ApJ*, 315, 209
- Prantzos, N., Hashimoto, M., & Nomoto, K. 1990, *A&A*, 234, 209 (P90)
- Raiteri, C.M., Busso, M., Gallino, R., Picchio, G., & Pulone, L. 1991, *ApJ*, 367, 228
- Raiteri, C.M., Gallino, R., & Busso, M. 1992, *ApJ*, 387, 263
- Raiteri, C.M., Gallino, R., Busso, M., Neuberger, D., & Käppeler, F. 1993, *ApJ*, 419, 207
- Schaller, G., Schaerer, D., Meynet, G., & Maeder, A. 1992 *A&AS*, 96, 269
- Schlegel, C. et al. 1994, *Phys. Rev.*, C50, 2198
- Schumann, M., Käppeler, F., Böttger, R., & Schölermann, H. 1997, *Nucl. Phys.*, A621, 274c
- Takahashi, K., & Yokoi, K. 1987, *Atomic Data Nucl. Data Tables* 36, 375.

- Thielemann F. K., Arnould M., & Truran J. W. 1986, in *Advances in Nuclear Astrophysics* ed. E. Vangioni-Flam, J. Audouze, M. Cassé, J.-P. Chièze, & J. Tran Thanh Van (Gif-sur-Yvette: Editions Frontieres), 525.
- Tuli J. K. 1995, *Nuclear Wallet Cards* (Upton, N.Y.: Brookhaven National Lab.)
- von Neumann-Cosel, P. et al. 1997, *Nucl. Phys.*, A621, 278c
- Weaver, T. A., & Woosley, W. E. 1993, *Phys. Rep.*, 227, 65
- Yokoi, K., & Takahashi, K. 1983, *Nature*, 305, 198

Table 1. Nuclei in the Network

Z	A_{min}	A_{max}	Z	A_{min}	A_{max}	Z	A_{min}	A_{max}	Z	A_{min}	A_{max}
1	2	2	22	42	51	43	97	101	64	152	161
2	3	4	23	44	52	44	96	105	65	156	162
3	7	7	24	46	55	45	103	105	66	156	165
4	7	7	25	48	56	46	102	111	67	162	166
5	8	11	26	50	61	47	107	111	68	162	171
6	11	15	27	51	61	48	106	117	69	168	172
7	13	16	28	54	65	49	103	117	70	168	177
8	15	19	29	56	66	50	112	127	71	174	180
9	17	20	30	59	71	51	121	127	72	174	183
10	20	23	31	69	72	52	120	131	73	179	182
11	21	24	32	70	77	53	125	131	74	180	188
12	23	27	33	75	77	54	124	137	75	184	189
13	25	28	34	74	83	55	131	137	76	184	195
14	26	31	35	79	83	56	130	139	77	190	195
15	27	34	36	78	87	57	137	140	78	190	199
16	30	37	37	85	88	58	136	144	79	195	200
17	33	38	38	84	91	59	141	143	80	196	205
18	34	41	39	89	91	60	142	151	81	203	206
19	37	42	40	90	97	61	144	151	82	204	209
20	38	49	41	93	97	62	144	155	83	209	210
21	40	49	42	92	101	63	151	157	84	210	254

Z is the proton number and A is the mass number.

Table 2. List of Stellar Models Studied

Case	$^{12}\text{C}(\alpha, \gamma)^{16}\text{O}$	$^{22}\text{Ne}(\alpha, n)^{25}\text{Mg}$	Star Mass ($M_{\odot}$)
A	CF88 $\times$ 1.7	CF88	15, 20, 25, 30
B	CF88 $\times$ 1.7	WT K94	15, 20, 25, 30
C	CF88	CF88	20
D	CFHZ85	CF88	20

Table 3. CHARACTERISTICS OF EVOLUTIONARY MODELS

stage	Evol. Time (yrs)	$\log(L/L_{\odot})$	$\log(T_{\text{eff}})$ (K)	T_c (10^8 K)	ρ_c (g cm^{-3})	M_{α} ($M_{\odot}$)	M_{co} ($M_{\odot}$)	M_{NeMg} ($M_{\odot}$)	$M_{\text{cc}}^{\text{max}}$ ($M_{\odot}$)
<u>15 $M_{\odot}$</u>									
1	0.00	4.274	4.484	0.340	6.13×10^0	0.00	0.00	0.00	5.58
2	$+1.06 \times 10^7$	4.617	4.405	0.628	6.30×10^1	2.30	0.00	0.00	0.00
3	$+3.30 \times 10^4$	4.641	4.186	1.30	1.31×10^3	2.69	0.00	0.00	2.17
4	$+1.68 \times 10^6$	4.733	3.560	3.12	6.31×10^3	3.90	2.07	0.00	0.00
5	$+3.14 \times 10^4$	4.816	3.550	5.46	1.10×10^5	3.91	2.12	0.00	0.51
6	$+8.12 \times 10^3$	4.849	3.548	9.81	5.54×10^6	3.91	2.21	1.12	0.00
7	$+4.85 \times 10^1$	4.845	3.548	12.11	1.35×10^7	3.91	2.24	1.47	0.00
<u>20 $M_{\odot}$</u>									
1	0.00	4.625	4.539	0.359	4.87×10^0	0.00	0.00	0.00	8.71
2	$+7.64 \times 10^6$	4.982	4.433	0.751	6.16×10^1	4.15	0.00	0.00	0.00
3	$+1.38 \times 10^3$	5.018	4.265	1.36	6.29×10^2	4.35	0.00	0.00	3.79
4	$+1.03 \times 10^6$	5.049	3.551	3.27	4.37×10^3	5.90	3.75	0.00	0.00
5	$+1.54 \times 10^4$	5.134	3.541	5.92	5.27×10^4	5.90	3.81	0.00	0.47
6	$+2.90 \times 10^3$	5.155	3.539	10.350	3.03×10^6	5.90	3.81	1.15	0.00
7	$+3.81 \times 10^1$	5.152	3.539	12.126	7.09×10^6	5.90	3.81	1.78	0.00
<u>25 $M_{\odot}$</u>									
1	0.00	4.877	4.575	0.372	4.06×10^0	0.00	0.00	0.00	12.26
2	$+6.19 \times 10^6$	5.236	4.395	0.972	1.07×10^2	6.11	0.00	0.00	0.00
3	$+5.40 \times 10^3$	5.258	4.267	1.40	3.91×10^2	6.36	0.00	0.00	5.65
4	$+7.65 \times 10^5$	5.279	3.079	3.61	4.33×10^3	7.95	5.49	0.00	0.00
5	$+9.40 \times 10^3$	5.341	3.535	6.24	3.79×10^4	7.95	5.52	0.00	0.30
6	$+1.30 \times 10^3$	5.358	3.533	12.07	1.63×10^6	7.95	5.54	1.56	0.00
7	$+5.25 \times 10^{-1}$	5.358	3.533	12.18	1.82×10^6	7.95	5.54	1.56	0.00

Table 3—Continued

stage	Evol. Time (yrs)	$\log(L/L_{\odot})$	$\log(T_{\text{eff}})$ (K)	T_c (10^8 K)	ρ_c (g cm $^{-3}$)	M_{α} ($M_{\odot}$)	M_{co} ($M_{\odot}$)	M_{NeMg} ($M_{\odot}$)	$M_{\text{cc}}^{\text{max}}$ ($M_{\odot}$)
				<u>30 $M_{\odot}$</u>					
1	0.00	5.070	4.601	0.381	3.54×10^0	0.00	0.00	0.00	16.06
2	$+5.32 \times 10^6$	5.428	4.263	1.36	2.45×10^2	8.51	0.00	0.00	0.00
3	$+5.00 \times 10^2$	5.428	4.233	1.44	2.97×10^2	8.51	0.00	0.00	7.65
4	$+6.17 \times 10^5$	5.439	3.557	3.43	2.94×10^3	10.06	7.55	0.00	0.00
5	$+6.52 \times 10^3$	5.439	3.550	6.34	2.71×10^4	10.06	7.59	0.00	0.00
6	$+7.90 \times 10^2$	5.505	3.550	13.11	1.21×10^6	10.06	7.59	2.02	0.00
7	$+0.00 \times 10^0$	5.505	3.550	13.11	1.21×10^6	10.06	7.59	2.02	0.00

^a $M_{\text{cc}}^{\text{max}}$ is the maximum convective core mass of that burning phase.

^b M_{α} is the mass size of helium core.

^c T_C is the central temperature.

^d ρ_C is the central density.

Stages 1 to 7 correspond to the evolutionary time of 1) Zero Age Main sequence, 2) central hydrogen exhaustion, 3) central helium ignition, 4) central helium exhaustion, 5) central carbon ignition, 6) central carbon exhaustion, and 7) central neon ignition, respectively.

Table 4. Production factors inside hydrogen exhausted core at the end of core hydrogen burning

isotope	Production factors			
	15 $M_{\odot}$	20 $M_{\odot}$	25 $M_{\odot}$	30 $M_{\odot}$
^1H	1.0×10^{-4}	7.6×10^{-4}	4.9×10^{-4}	9.3×10^{-4}
^3He	3.4×10^{-12}	8.3×10^{-11}	4.2×10^{-11}	1.1×10^{-9}
^4He	3.6×10^0	3.6×10^0	3.6×10^0	3.6×10^0
^7Li	2.4×10^{-13}	2.1×10^{-12}	1.3×10^{-12}	2.6×10^{-12}
^{12}C	8.9×10^{-2}	9.3×10^{-2}	9.8×10^{-2}	1.0×10^{-2}
^{13}C	2.3×10^0	2.4×10^0	2.5×10^0	2.6×10^0
^{14}N	1.1×10^1	1.1×10^1	1.1×10^1	1.1×10^1
^{15}N	9.0×10^{-2}	9.3×10^{-2}	9.1×10^{-2}	9.2×10^{-2}
^{16}O	3.2×10^{-2}	3.0×10^{-2}	2.9×10^{-2}	2.7×10^{-2}
^{17}O	9.7×10^{-2}	9.1×10^{-2}	9.0×10^{-2}	8.7×10^{-2}
^{18}O	5.7×10^{-6}	5.4×10^{-6}	4.8×10^{-6}	4.5×10^{-6}
^{19}F	2.6×10^{-5}	2.6×10^{-5}	3.0×10^{-5}	3.1×10^{-5}
^{20}Ne	9.9×10^{-1}	9.9×10^{-1}	9.8×10^{-1}	9.8×10^{-1}
^{21}Ne	1.9×10^0	7.0×10^{-1}	2.1×10^{-1}	1.1×10^{-1}
^{22}Ne	2.6×10^{-2}	1.7×10^{-2}	9.8×10^{-3}	6.1×10^{-3}
^{23}Na	5.5×10^0	5.9×10^0	6.0×10^0	6.1×10^0
^{25}Mg	6.0×10^{-2}	4.4×10^{-2}	3.7×10^{-2}	3.2×10^{-2}
^{26}Mg	1.8×10^0	1.8×10^0	1.8×10^0	1.8×10^0
^{138}La	9.5×10^{-1}	9.6×10^{-1}	9.6×10^{-1}	9.6×10^{-1}
^{163}Dy	6.4×10^{-1}	7.9×10^{-1}	7.0×10^{-1}	6.9×10^{-1}
^{187}Re	3.6×10^{-2}	1.0×10^{-2}	1.6×10^{-2}	1.6×10^{-2}
^{187}Os	5.2×10^0	5.3×10^0	5.3×10^0	5.3×10^0

Table 5. Central ^{12}C and ^{16}O Mass Fractions at Core Helium Exhaustion

Model	$X(^{12}\text{C})$	$X(^{16}\text{O})$
15 $M_{\odot}$, case A	0.251	0.723
15 $M_{\odot}$, case B	0.219	0.753
20 $M_{\odot}$, case A	0.231	0.741
20 $M_{\odot}$, case B	0.218	0.752
20 $M_{\odot}$, case C	0.335	0.637
20 $M_{\odot}$, case D	0.115	0.856
25 $M_{\odot}$, case A	0.220	0.750
25 $M_{\odot}$, case B	0.211	0.757
30 $M_{\odot}$, case A	0.226	0.741
30 $M_{\odot}$, case B	0.217	0.750

Table 6. Comparing Stellar Model Parameters Relevant to s-process: Maximum Convective Core Mass & Duration of Helium Burning

Stellar Model	15 $M_{\odot}$		20 $M_{\odot}$		25 $M_{\odot}$		30 $M_{\odot}$	
	M_{cc}^{max} ($M_{\odot}$)	Δt_{He} (10^6 yrs)	M_{cc}^{max} ($M_{\odot}$)	Δt_{He} (10^6 yrs)	M_{cc}^{max} ($M_{\odot}$)	Δt_{He} (10^6 yrs)	M_{cc}^{max} ($M_{\odot}$)	Δt_{He} (10^6 yrs)
FRANEC (K94)	2.6	1.19	4.4	0.781	6.5	0.620	8.6	0.525
Schaller et al. (1992)	2.80	1.12	4.83	0.79	7.13	0.63	-	-
Chieffi et al. (1998)	-	-	-	-	5.34	0.68	-	-
Prantzos et al. (1990)	1.98	-	3.70	1.04	5.66	0.795	7.50	-
Present Work $1.7 \times {}^{12}\text{C}(\alpha, \gamma)$ CF88	2.17	1.68	3.79	1.03	5.65	0.762	7.65	0.617
Present Work $1.0 \times {}^{12}\text{C}(\alpha, \gamma)$ CF88	-	-	3.71	1.01	-	-	-	-
Present Work $1.0 \times {}^{12}\text{C}(\alpha, \gamma)$ CFHZ85	-	-	3.85	1.08	-	-	-	-

M_{cc}^{max} is the maximum mass radius of convective core

Δt_{He} is the duration of core helium burning

Parameters of FRANEC results are obtained from figure 10 of K94 in which ${}^{12}\text{C}(\alpha, \gamma){}^{16}\text{O}$ rate of CFHZ85 were used

Schaller et al. (1992) use overshooting method during the helium convective phase

Chieffi et al. (1998) use ${}^{12}\text{C}(\alpha, \gamma){}^{16}\text{O}$ rate of CFHZ85

Parameters of Prantzos et al. (1990)'s models are obtained from their Table 1 and Nomoto & Hashimoto (1988)

Table 7. Comparison of s-Processing in Massive Stars during Core Helium Burning among Authors when $^{22}\text{Ne}(\alpha, n)^{25}\text{Mg}$ rate of CF88 is used

Author	τ_c	n_c^a	$\langle \tau \rangle$ (mb^{-1}) ^b	n_n^{max} ($\times 10^5 \text{ cm}^{-3}$) ^c	X_{22} ($\times 10^{-2}$) ^d	$X_{80}/X_{80\odot}^e$
<u>15 $M_\odot$</u>						
P90	–	3.20	0.19	–	1.23	72
K94	–	1.85	0.09	2.05	1.65	21
A	4.00	3.38	0.10	2.27	1.33	117
<u>20 $M_\odot$</u>						
P90	–	4.90	0.27	–	0.92	342
K94	–	3.66	0.15	5.06	1.32	116
A	5.93	5.48	0.16	3.50	1.04	598
C	6.69	6.47	0.17	2.10	0.81	1010
D	6.02	5.85	0.16	3.80	0.94	719
<u>25 $M_\odot$</u>						
P90	–	6.20	0.33	–	0.68	786
K94	–	5.41	0.20	6.62	1.00	475
A	7.15	6.70	0.22	4.24	0.76	1100
<u>30 $M_\odot$</u>						
P90	–	7.10	0.37	–	0.53	1177
K94	–	6.55	0.23	6.74	0.79	933
A	8.09	7.36	0.22	4.44	0.65	1368

P90 is the results from P90 with $^{12}\text{C}(\alpha, \gamma)^{16}\text{O}$ rate of CF88.

K94 is the results from K94 with $^{12}\text{C}(\alpha, \gamma)^{16}\text{O}$ rate of CFHZ85 (their Table 5).

A, C, & D are the results of the present work.

τ_c is the central neutron exposure according to Eq. (3) in the text.

^aNumber of neutrons captured per iron seed averaged over the maximum convective core mass.

^bMean Neutron exposure at 30 keV, averaged over the core as τ_c .

^cMaximum of the mean neutron density.

^dFinal ^{22}Ne mass fraction.

^eFinal ^{80}Kr production factor averaged over the maximum convective core mass.

Table 8. Comparison of s-Processing in Massive Stars during Core Helium Burning among Authors when the tentative $^{22}\text{Ne}(\alpha, n)^{25}\text{Mg}$ rate of K94 is used

Author	τ_c	n_c^a	$\langle \tau \rangle$ (mb^{-1}) ^b	n_n^{max} ($\times 10^5 \text{ cm}^{-3}$) ^c	X_{22} ($\times 10^{-2}$) ^d	$X_{80}/X_{80\odot}^e$
				<u>15 $M_\odot$</u>		
K94	–	6.61	0.23	3.25	1.10	967
B	9.09	11.1	0.28	2.41	0.40	3883
				<u>20 $M_\odot$</u>		
K94	–	10.7	0.31	4.93	0.50	3895
B	10.42	13.8	0.28	2.16	0.19	4918
				<u>25 $M_\odot$</u>		
K94	–	12.9	0.35	4.29	0.17	5771
B	11.74	14.0	0.31	2.02	0.10	5332
				<u>30 $M_\odot$</u>		
K94	–	13.6	0.36	3.74	0.07	6379
B	14.50	14.4	0.33	2.28	0.07	5498

All entries have their meanings as in Table 5.

B are the results of the present work with $1.7 \times ^{12}\text{C}(\alpha, \gamma)^{16}\text{O}$ rate of CF88.

Table 9. Overproduction factors $f=X/X_{\odot}$ within the maximum convective core for nuclei in the network with $Z \leq 40$ and $f > 0.05$ and some nuclei with $Z > 40$ and $f > 10$

isotope	Proton Number	Mass Number	f using $^{22}\text{Ne}(\alpha, n)^{25}\text{Mg}$ of CF88				f using $^{22}\text{Ne}(\alpha, n)^{25}\text{Mg}$ of WT K94			
			15 $M_{\odot}$	20 $M_{\odot}$	25 $M_{\odot}$	30 $M_{\odot}$	15 $M_{\odot}$	20 $M_{\odot}$	25 $M_{\odot}$	30 $M_{\odot}$
c12	6	12	82.3	75.3	72.4	75.4	71.6	71.4	70.0	72.1
o16	8	16	74.0	75.6	76.6	76.7	76.9	77.4	77.8	77.6
o18	8	18	5.5	2.0	4.8	0.0	7.1	4.1	0.2	0.0
ne20	10	20	1.2	1.9	2.9	3.7	1.3	2.1	3.1	4.1
ne21	10	21	3.4	3.6	3.9	4.2	5.0	4.2	3.6	3.2
ne22	10	22	107.2	79.2	60.0	51.6	39.0	14.5	6.9	3.9
na23	11	23	7.6	8.0	8.1	8.2	7.7	7.4	7.3	7.3
mg24	12	24	0.9	0.9	0.9	1.0	0.8	0.8	0.9	1.0
mg25	12	25	44.4	68.6	81.5	89.0	125.5	146.7	153.5	157.3
mg26	12	26	41.9	78.1	101.0	114.3	103.9	133.7	142.6	147.3
al27	13	27	0.7	0.8	0.9	0.9	0.9	1.0	1.1	1.1
si28	14	28	0.9	0.8	0.8	0.8	0.8	0.7	0.7	0.7
si29	14	29	2.1	2.3	2.3	2.4	2.1	2.2	2.2	2.2
si30	14	30	1.9	2.4	2.6	2.7	2.4	2.6	2.6	2.7
p31	15	31	7.0	9.9	11.4	12.3	15.8	17.5	18.2	18.6
s32	16	32	0.6	0.5	0.5	0.4	0.4	0.4	0.4	0.3
s33	16	33	30.3	31.6	31.5	31.7	27.8	26.9	26.5	26.6
s34	16	34	4.0	5.6	6.3	6.7	7.5	7.9	8.1	8.2
s36	16	36	1.1	1.2	1.9	2.9	1.6	1.7	2.0	2.2
cl35	17	35	4.6	7.5	9.0	9.8	12.2	13.6	14.0	14.3
cl37	17	37	59.2	69.5	74.6	78.1	89.8	98.5	101.4	103.4
ar36	18	36	0.3	0.2	0.2	0.1	0.1	0.1	0.1	0.1
ar38	18	38	1.3	1.6	1.8	1.9	2.3	2.6	2.6	2.7
ar40	18	40	3.2	7.2	8.5	6.9	13.2	22.9	20.4	21.5
k39	19	39	1.2	1.4	1.5	1.6	2.0	2.2	2.3	2.3
k40	19	40	281.9	327.2	359.2	379.1	470.5	522.6	545.5	556.4
k41	19	41	15.2	13.3	12.9	12.8	13.7	14.5	14.9	15.1
ca40	20	40	0.4	0.2	0.2	0.2	0.1	0.1	0.1	0.1
ca42	20	42	36.5	31.8	29.1	28.2	22.7	21.4	21.0	20.9
ca43	20	43	46.7	43.8	40.9	40.1	30.8	29.1	28.6	28.5
ca44	20	44	10.4	13.9	14.6	15.1	14.5	13.9	13.8	13.8
ca46	20	46	0.7	1.8	1.7	1.1	1.8	2.7	2.4	2.7
ca48	20	48	0.9	0.8	0.8	0.8	0.7	0.7	0.7	0.7
sc45	21	45	31.6	45.4	49.1	51.4	48.8	47.4	47.0	47.2
ti46	22	46	14.6	25.2	28.9	30.9	34.2	33.5	33.3	33.5
ti47	22	47	4.6	9.0	10.8	11.8	12.9	13.0	13.0	13.1
ti48	22	48	0.6	1.4	1.8	2.0	2.5	2.5	2.6	2.6
ti49	22	49	8.0	18.9	25.7	29.3	43.4	46.5	47.7	48.5
ti50	22	50	18.0	32.7	44.9	51.7	104.9	130.4	139.1	143.2

Table 9—Continued

isotope	Proton Number	Mass Number	f using $^{22}\text{Ne}(\alpha, n)^{25}\text{Mg}$ of CF88				f using $^{22}\text{Ne}(\alpha, n)^{25}\text{Mg}$ of WT K94			
			15 $M_{\odot}$	20 $M_{\odot}$	25 $M_{\odot}$	30 $M_{\odot}$	15 $M_{\odot}$	20 $M_{\odot}$	25 $M_{\odot}$	30 $M_{\odot}$
v51	23	51	0.5	0.8	1.2	1.4	2.6	3.4	3.7	3.8
cr52	24	52	0.4	0.3	0.3	0.3	0.3	0.4	0.4	0.4
cr53	24	53	0.4	0.3	0.3	0.3	0.3	0.3	0.4	0.4
cr54	24	54	16.3	15.6	14.9	14.8	12.7	12.9	12.9	13.1
mn55	25	55	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.1
fe56	26	56	0.3	0.2	0.1	0.1	0.1	0.1	0.1	0.0
fe57	26	57	3.9	2.4	1.9	1.6	0.9	0.8	0.7	0.6
fe58	26	58	101.0	81.4	70.2	65.6	41.2	33.8	31.3	30.3
co59	27	59	34.6	31.2	27.9	26.6	16.7	14.0	13.0	12.6
ni60	28	60	7.5	8.0	7.6	7.4	5.0	4.3	4.0	3.9
ni61	28	61	52.2	59.4	57.4	57.0	39.1	33.4	31.4	30.9
ni62	28	62	30.8	40.6	41.4	42.2	30.8	26.8	25.5	25.2
ni64	28	64	117.0	226.0	269.9	293.4	314.6	302.5	298.5	300.4
cu63	29	63	57.5	80.5	82.8	84.7	65.8	57.3	54.1	53.6
cu65	29	65	99.5	193.1	231.2	251.1	265.8	252.5	249.3	251.0
zn64	30	64	14.2	22.4	24.2	25.4	20.4	18.3	17.6	17.6
zn66	30	66	44.8	101.9	130.4	145.6	171.4	169.2	169.5	171.4
zn67	30	67	62.1	147.6	192.1	216.3	261.5	260.5	261.8	265.0
zn68	30	68	54.2	158.4	225.1	262.3	402.4	429.0	440.8	449.0
zn70	30	70	3.7	13.5	21.6	26.0	58.9	70.3	74.1	75.8
ga69	31	69	63.5	194.5	281.4	330.0	525.2	566.1	583.8	595.2
ga71	31	71	76.3	268.0	409.3	489.2	868.2	971.1	1012.3	1034.8
ge70	32	70	81.6	269.6	401.5	475.4	811.8	891.3	924.2	943.4
ge72	32	72	54.2	210.1	334.3	404.0	804.0	924.8	971.6	995.3
ge73	32	73	33.0	133.0	215.5	262.8	519.9	609.2	643.7	660.6
ge74	32	74	26.7	117.7	197.9	242.9	554.8	666.3	707.8	727.0
ge76	32	76	0.1	0.1	0.1	0.0	0.2	0.2	0.1	0.1
as75	33	75	19.4	87.3	148.3	182.8	419.7	508.6	542.0	557.4
se76	34	76	55.8	259.7	447.5	552.5	1342.2	1640.1	1750.0	1799.7
se77	34	77	23.0	109.0	189.5	234.8	576.5	709.8	759.3	781.6
se78	34	78	28.0	141.0	252.6	314.9	840.0	1059.3	1139.0	1173.9
se80	34	80	2.3	13.5	24.7	31.2	95.2	129.4	135.4	139.4
br79	35	79	8.1	41.1	74.1	92.4	248.1	314.0	337.0	347.0
br81	35	81	3.7	28.0	45.0	36.5	154.3	221.3	210.0	194.2
kr80	36	80	117.3	606.3	1100.5	1374.6	3883.1	4946.6	5337.9	5509.6
kr82	36	82	54.9	304.5	578.4	731.0	2329.6	3099.9	3368.3	3485.6
kr83	36	83	15.4	88.1	171.1	217.5	697.7	950.2	1039.2	1076.6
kr84	36	84	12.1	71.8	147.8	189.4	722.1	1044.9	1158.2	1204.2
kr86	36	86	1.0	5.7	8.0	6.6	49.3	143.3	116.5	132.0
rb85	37	85	8.1	48.3	100.2	127.3	508.6	741.6	815.2	845.8
rb87	37	87	0.7	2.5	3.7	2.7	24.2	71.8	65.3	74.9

Table 9—Continued

isotope	Proton Number	Mass Number	f using $^{22}\text{Ne}(\alpha, n)^{25}\text{Mg}$ of CF88				f using $^{22}\text{Ne}(\alpha, n)^{25}\text{Mg}$ of WT K94			
			15 $M_{\odot}$	20 $M_{\odot}$	25 $M_{\odot}$	30 $M_{\odot}$	15 $M_{\odot}$	20 $M_{\odot}$	25 $M_{\odot}$	30 $M_{\odot}$
sr86	38	86	46.8	273.9	587.6	764.1	3369.9	5121.8	5758.8	6016.6
sr87	38	87	40.2	230.6	507.9	662.0	3152.6	4932.2	5587.7	5841.2
sr88	38	88	13.2	57.5	127.7	167.6	1085.8	2007.4	2311.3	2438.9
y89	39	89	9.8	35.5	76.0	99.5	756.2	1550.1	1794.5	1903.2
zr90	40	90	4.3	13.6	27.3	35.1	298.6	673.8	781.8	833.1
zr91	40	91	5.1	15.5	30.4	39.0	332.7	793.9	924.5	987.8
zr92	40	92	4.4	12.4	22.9	28.8	252.4	641.5	746.1	800.0
zr94	40	94	3.4	8.6	14.7	18.1	151.4	422.7	490.2	528.9
zr96	40	96	0.2	0.1	0.1	0.1	0.1	0.5	0.3	0.3
mo96	42	96	3.1	7.7	12.8	15.6	124.9	361.8	420.2	454.7
mo98	42	98	2.2	5.2	8.5	10.2	78.8	233.7	270.5	293.2
ru100	44	100	2.7	6.0	9.7	11.7	87.6	263.8	305.0	331.0
pd104	46	104	2.6	5.6	8.9	10.6	74.9	230.5	266.2	289.6
gd152	64	152	22.4	35.3	43.1	47.3	64.8	75.5	79.6	81.9
ta180	73	180	1.0	4.5	9.4	7.4	3.3	11.1	16.6	15.3




Fig. 1.— Evolution of the central temperature vs. central density for the four case A stellar models. The temperature scale, $\tilde{T}_C$ is $\approx 2 \times 10^7 \rho^{1/3}$. All four star models have been evolved up to central neon ignition.




Fig. 2.— The convective helium-burning core mass in the four case A star models versus the fraction of the helium-burning time (upper panel) and the mass fraction of ^4He (lower panel). Starting at $X(^4\text{He}) = 0.1$, the edges of the convective cores have been finely resolved to prevent a sudden injection of a large supply of ^4He into the cores.




Fig. 3.— Several s-process diagnostics for two $25 M_{\odot}$ stellar models. The case A model is shown as the solid curve while the case B model is shown as the dashed line. The diagnostics are the following: a) τ_n is the central neutron exposure, as computed from Eq. (3). It is the neutron exposure that would have been seen by a nucleus that stayed in the center of the star throughout the burning (τ_n). τ_{54} is the neutron exposure computed from the decline of the ^{54}Fe abundance due to neutron captures (see Eq. [5]). The latter better indicates the average neutron exposure of the core helium-burning s-process because it reflects the dilution by the convective mixing. It also shows that the bulk of the s-processing occurs mainly in the range $X(^4\text{He})_c \simeq 0.01 - 0.001$. b) Overabundances of key isotopes. Note that the production factor of ^{22}Ne is multiplied by 1/100 in order that it not overlap in the figure that of ^{25}Mg . c) The number of neutrons captured per iron-seed nucleus according to Eq. (2). d) The central neutron density of the star during core helium burning. All quantities are plotted against central helium mass fraction. A low-lying $^{22}\text{Ne}(\alpha, n)^{25}\text{Mg}$ resonance (case B) would dramatically increase the robustness of the core helium-burning s-process.




Fig. 4.— The central neutron density as function of central helium mass fraction or time for the four case A (top panel) and the four case B (lower panel) stellar models.




Fig. 5.— The neutron exposure τ_n as a function of interior mass for all eight case A (solid curve) and B (dashed curve) stellar models through core helium burning. τ_n is the neutron exposure a nucleus would experience were it to remain at the particular interior mass. This figures clearly demonstrates that most neutron captures occur in the very center of the star. Nuclei are convected down into the center of the star, capture neutrons, and then mix back out into the core.




Fig. 6.— Similar to Figure 3 but for the case A (solid curve), case C (long-dashed curve), and case D (short-dashed curve) $20 M_{\odot}$ stellar models. The lowest $^{12}\text{C}(\alpha, \gamma)^{16}\text{O}$ reaction rate gives the largest degree of s-processing. The results of cases A and D agree to within $\sim 10 - 20\%$.




Fig. 7.— Overabundances of heavy nuclei (mass number $A \geq 12$) averaged over the convection helium-burning core for the four case A stellar models. The primary nucleosynthesis production process for each isotope is indicated by the symbol type.




Fig. 8.— Overabundances of heavy nuclei (mass number $A \geq 12$) averaged over the convection helium-burning core for the $25 M_{\odot}$ stellar model with 10% contribution to the reaction rate for $^{22}\text{Ne}(\alpha, n)^{25}\text{Mg}$ from a low-lying resonance.




Fig. 9.— Several s-process diagnostics for the 25 $M_{\odot}$ case A stellar model at central neon ignition. The upper panel shows the two measures of the neutron exposure. Of particular interest is the run of τ_n against interior mass. The dashed line shows the conditions at the end of core helium burning, while the solid line shows the conditions at the end of core neon burning. Most s-processing has occurred in the center of the star, but the contribution from shell carbon burning and shell helium burning are evident at $1.6 M_{\odot}$ and about $5.4 M_{\odot}$, respectively. The middle panel shows the number of neutrons captured per iron seed nucleus and the overabundance of ^{80}Kr at the end of core helium burning (dashed line) and neon burning (solid line). Various convective burning phases after central helium burning modify the number of neutrons captured per seed nucleus. Interestingly, shell carbon s-processing has destroyed a significant fraction of the ^{80}Kr produced in earlier convection core helium burning. The bottom panel shows mass fractions at the end of core neon burning of some nuclei to indicate the nuclear burning region.